

# Vaccination Schedule 2026

DISEASE	VACCINE	BIRTH	6 WEEKS	10 WEEKS	14 WEEKS	6 MONTHS	9 MONTHS	12 MONTHS	15 MONTHS	18 MONTHS	6 YEARS	9 YEARS	12 YEARS	
Tuberculosis (TB)	Bacillus Calmette-Guérin (BCG)	★												
Polio	Bivalent Oral Polio Vaccine (bOPV)	★	★											
Diphtheria   Tetanus   Polio   Pertussis   Haemophilus influenzae   Hepatitis B	Hexaxim		★	★	★					★				
	Infanrix <sup>®</sup> .hexa		★	★	★	The Expanded Program on Immunisation schedule (at 6, 10, 14 weeks of age) may only be used if a dose of hepatitis B vaccine has been given at birth.								
Pneumococcal	Prevenar 13		★	★	★			★	★					
	Vaxneuvance <sup>®</sup>		★					★	★					
	Synflorix		★		★		★		Can be administered as of 6 WEEKS of age. Official recommendations should be followed when immunising with Synflorix.					
	PCV-10 Cipla		★		★		★							
Rotavirus	Rotarix		★		★	The vaccination course consists of two doses. The first dose may be administered from the age of 6 WEEKS. There should be an interval of at least 4 weeks between doses. The vaccination course should be completed by the age of 24 WEEKS.								
	RotaTeq <sup>®</sup>		★	★	★									
Measles   Mumps   Rubella	Priorix <sup>®</sup>							★			★			
	Omzyta <sup>®</sup>							★			★			
Measles   Rubella	MR SII					★		★						
Measles	Cipla-Measles					★								
Chickenpox	Varilrix <sup>®</sup>								Children from 9 MONTHS up to and including 12 YEARS of age should receive 2 doses (second dose at least 6 weeks after the first dose)					
	Onvara <sup>®</sup>								★					
Meningococcal	Bexsero (Men B)			★	★			★	★					
	Menactra (Men A, C, W, Y)							★	★	For individuals aged 2 - 55 years, administer 1 dose.				
Hepatitis A	Havrix <sup>®</sup> Junior							★		★				
	Avaxim 80							★		★				
Tetanus   Diphtheria   Pertussis with or without Polio	Boostrix <sup>®</sup> tetra										★ FROM 3 YEARS ONWARDS			
	Boostrix <sup>®</sup>										★ FROM 4 YEARS ONWARDS			
	Tetraxim										★			
	Adacel Quadra										★		★	
	Tdap										★		★	
Human Papilloma Virus (HPV)	Cervarix <sup>®</sup>					Cervarix: For ages 9-14 years, administer 2 doses 6 months apart. As part of the state programme, girls and boys in public schools aged 9-14 years receive Cervarix in 2 doses, 6 months apart.						★		
	Gardasil <sup>®</sup>					Gardasil: For ages 9-13 years, administer 2 doses 6 months apart.						★		
	Gardasil <sup>®</sup> 9					Gardasil 9: For ages 9-14 years, administer 2 doses 5-13 months apart.						★		
Influenza	Influvac					Recommended from 6 months of age, especially for children in creches, or with chronic illnesses or respiratory problems. Must be administered before the influenza season in May.								
	Vaxigrip Tetra													
Respiratory Syncytial Virus (RSV)	Synagis <sup>®</sup>					RSV Prophylaxis is recommended for high-risk infants to prevent serious lower respiratory tract infections (LRTI) caused by RSV. Administer from January to May.								
Pneumococcal	Pneumovax 23					Children must receive at least one dose of the Pneumococcal Conjugate Vaccine (PCV) before receiving Pneumovax 23. Pneumovax 23 is only administered to children older than 2 years who are immune-compromised or at high risk of pneumococcal infection. Two doses of Pneumovax 23 are recommended, with the second dose given 3-8 years after the first dose.								
Hepatitis B	Engerix <sup>®</sup> -B										The vaccine can be used in subjects from 11 years up to and including 15 years of age as a 2-dose schedule.			
	Euvax		★											
	SII Hepatitis B Vaccine	★				This vaccine is administered to babies who are exposed to the hepatitis B surface antigen (HbsAg) at birth.								



Scan this QR code to access the full prescribing information for Aspen/ GSK vaccinations.



**OPTIONAL VACCINES**  
Some of the vaccines in this schedule are only available at private clinics.

**EXPANDED PROGRAM ON IMMUNISATION (EPI) VACCINES**  
These vaccines are available free of charge from government-supplied clinics.